

By-O-reg+ 360 Grow/Finish pigs Information

Field usage (> 9,000 pigs) comparison vs Ionophore

- 8.6 days **quicker to market** when fed BOR+ 360
- 1% ↓ mortality (in absolute terms)
- 2.8% ↓ late mortality due to HBS (in absolute terms)
- 2.2% ↓ FCR
- 3.8% ↑ bodyweight
- ↓ medicine costs by \$2.84/pig
- 41% ↓ salivary cortisol levels

Findings from University trial vs Ionophore

- 1.3% ↑ body weight gain
- 5.2% ↓ FCR
- 2.1% ↑ hot carcass weight
- 1.9% ↑ dressing %
- 5.0% ↑ loin area
- 7.5% ↓ cooking loss
- 3.6% ↑ backfat (10th & last rib)
- Maintains feed intake & ↑ body weight gain under heat stress vs Ionophore where feed intake ↑ with ↓ body weight gain

Findings from University trials vs negative control

- 4% ↑ body weight gain
- No difference FCR
- 5% ↑ hot carcass weight
- 1.9% ↑ dressing %
- 7.6% ↑ loin area
- 4.7% ↓ cooking loss
- 6.5% ↑ backfat (10th & last rib)



*These data come from university studies and on-farm trials.