

START-N-GROW

22|20 CALF MILK REPLACER

PRODUCT CODE(S): 59885

PRODUCT DESCRIPTION: Provides Essential Nutrition & Advance Support from Day 2 through Weaning, with 22% Protein and 20% Fat to Support Optimal Growth

STORAGE

Store at room temperature in a dry place. Use mixed product immediately – do not store.

AVAILABLE SIZES

50 lb. Bag

MIXING DIRECTIONS

- 1 - Start with 1 Quart of Warm Water (120°F)
- 2 - Add 10 oz of Milk Replacer Powder
- 3 - Mix Thoroughly Until Smooth
- 4 - Add Warm Water Until Full Mixture Equals 2 qts
- 5 - Pour & Feed (102°F)

GUARANTEED ANALYSIS

Crude Protein, minimum.....	22.0%
Lysine, minimum.....	2.4%
Crude Fat, minimum.....	20.0%
Crude Fiber, maximum.....	0.5%
Calcium (Ca).....min...0.9%.....max...1.0%	
Phosphorus (P), minimum.....	0.6%
Selenium (Se), minimum	0.3 ppm
Vitamin A, minimum	16,000 IU/lb
Vitamin E, minimum	160 IU/lb

INGREDIENTS

Dried Whey, Dried Whey Protein Concentrate, Animal Fat (preserved with BHA & BHT), Condensed Whey, Vegetable Fats, Monoglycerides of Edible Oils, Silicon Dioxide, Dried Skim Milk, L-Lysine Monohydrochloride, Dicalcium Phosphate, L-Threonine, DL-Methionine, Calcium Carbonate, Natural and Artificial Flavors, Precipitated Silica, Dextrose, Lecithin, Inulin, Oregano Oil, Yeast Extract, Cinnamon Bark Oil, Hydrated Sodium Calcium Aluminosilicate, Ascorbic Acid, Ferrous Sulfate, Manganese Sulfate, Zinc Sulfate, Iron Proteinate, Manganese Proteinate, Selenium Yeast, Brewers Dried Yeast, Vitamin D3 Supplement, Sodium Selenite, D-Calcium Pantothenate, Copper Proteinate, Niacinamide, Maltodextrin, Biotin, Calcium Iodate, Riboflavin, Folic Acid, Vitamin B12 Supplement, Pyridoxine Hydrochloride, Thiamine Mononitrate, Menadione Sodium Bisulfite Complex (source of Vitamin K3), Cobalt Sulfate

FEEDING DIRECTIONS

AGE OF CALF	MILK REPLACER	FEEDINGS PER DAY
Day 1	Colostrum	
Day 2 - Week 8	2 Quarts	2 Feedings
For example, an average 2 day old calf would receive: 2 qt of Milk Replacer per feeding Fed 2 times per day		

Mix START-N-GROW Calf Milk Replacer into warm (120°F) water. Stir until the powder is completely dissolved. Only mix enough powder for one feeding at a time.

